



# Very nilpotent basis and n-tuples in Borel subalgebras

Michael Bulois

## ► To cite this version:

Michael Bulois. Very nilpotent basis and n-tuples in Borel subalgebras. *Comptes Rendus. Mathématique*, 2011, 349 (3-4), pp.149-152. 10.1016/j.crma.2010.12.005 . hal-00660382

**HAL Id: hal-00660382**

**<https://hal.science/hal-00660382>**

Submitted on 16 Jan 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Bases fortement nilpotentes et $n$ -uplets des algèbres de Borel

## Very nilpotent basis and $n$ -tuples in Borel subalgebras

MICHAËL BULOIS \*

### Résumé

Une base (d'espace vectoriel)  $B$  d'une algèbre de Lie est dite fortement nilpotente si tous les crochets itérés des éléments de  $B$  sont nilpotents. Dans cette note, on démontre une version améliorée du théorème d'Engel. On montre qu'une algèbre de Lie admet une base fortement nilpotente si et seulement si c'est une algèbre nilpotente. Lorsque  $\mathfrak{g}$  est une algèbre de Lie semi-simple, ceci nous permet de définir un idéal de  $S((\mathfrak{g}^n)^*)^G$  dont l'ensemble algébrique associé dans  $\mathfrak{g}^n$  est l'ensemble des  $n$ -uplets vivants dans une même sous-algèbre de Borel.

### Abstract

A (vector space) basis  $B$  of a Lie algebra is said to be very nilpotent if all the iterated brackets of elements of  $B$  are nilpotent. In this note, we prove a refinement of Engel's Theorem. We show that a Lie algebra has a very nilpotent basis if and only if it is a nilpotent Lie algebra. When  $\mathfrak{g}$  is a semisimple Lie algebra, this allows us to define an ideal of  $S((\mathfrak{g}^n)^*)^G$  whose associated algebraic set in  $\mathfrak{g}^n$  is the set of  $n$ -tuples lying in a same Borel subalgebra.

Rubriques: Algèbres de Lie/ Géométrie algébrique

## 1 Introduction and Notation

Let  $\mathfrak{g}$  be a Lie algebra defined over an algebraically closed field  $\mathbb{k}$  of characteristic 0. The adjoint action of  $z \in \mathfrak{g}$ :  $x \mapsto [z, x]$  is denoted by  $\text{ad}_z \in \mathfrak{gl}(\mathfrak{g})$ . If  $h \in \mathfrak{g}$ , we denote the centralizer of  $h$  in  $\mathfrak{g}$  by  $\mathfrak{g}^h$ . When  $V$  is a vector space over  $\mathbb{k}$ , and  $A \subset V$ ,  $\langle A \rangle$  stands for the linear subspace spanned by  $A$ . The symmetric algebra on  $V$  is denoted by  $S(V)$ . Any subset  $J \subset S(V^*)$  defines an algebraic subset  $\mathcal{V}(J) := \{x \in V \mid f(x) = 0, \forall f \in J\}$ .

For  $n \in \mathbb{N}^*$ , let  $\mathcal{I}_n$  be the set of morphisms of varieties  $\mathfrak{g}^n \rightarrow \mathfrak{g}$  defined by induction as follows:

- For  $i \in \llbracket 1, n \rrbracket$ ,  $((y_1, \dots, y_n) \mapsto y_i) \in \mathcal{I}_n$ .
- If  $f, g \in \mathcal{I}_n$ , then  $[f, g] := ((y_1, \dots, y_n) \mapsto [f(y_1, \dots, y_n), g(y_1, \dots, y_n)]) \in \mathcal{I}_n$ .

In particular,  $\mathcal{I}(y_1, \dots, y_n) := \{f(y_1, \dots, y_n) \mid f \in \mathcal{I}_n\}$  is the set of iterated brackets in  $y_1, \dots, y_n$ . One defines the depth map on  $\mathcal{I}_n$  by induction:

$$\text{dep}((y_1, \dots, y_n) \mapsto y_i) = 1, \quad \text{dep}[f, g] = \max\{\text{dep } f, \text{dep } g\} + 1.$$

We say that  $(y_1, \dots, y_n)$  is a *very nilpotent* basis of  $\mathfrak{g}$  if the following two conditions hold:

- $(y_1, \dots, y_n)$  is a basis of the vector space  $\mathfrak{g}$ ,
- $\text{ad}_z$  is nilpotent in  $\mathfrak{gl}(\mathfrak{g})$  for any  $z \in \mathcal{I}(y_1, \dots, y_n)$ .

The key result of this note is:

**Proposition 1.**  *$\mathfrak{g}$  has a very nilpotent basis if and only if  $\mathfrak{g}$  is nilpotent.*

Proposition 1 can be seen as a refinement of Engel's theorem (see, e.g. [TY, 19.3.6]). Its proof is rather technical and is given in section 2.

Assume now that  $\mathfrak{g}$  is semisimple. Let  $G$  be the algebraic adjoint group of  $\mathfrak{g}$  and let  $p_1, \dots, p_d$  be algebraically independant homogeneous generators of  $S(\mathfrak{g}^*)^G$ , the set of  $G$ -invariant elements of  $S(\mathfrak{g}^*)$ . It is well known that  $\mathcal{V}(p_1, \dots, p_d)$  is the nilpotent cone of  $\mathfrak{g}$  (see, e.g. [TY, §31]). Let  $J_0$  be the ideal of  $S((\mathfrak{g}^n)^*) = \bigotimes_n S(\mathfrak{g}^*)$  generated by the polynomials  $p_i \circ f$  where  $f \in \mathcal{I}_n$ . We define  $J$  in the same way, adding the constraint  $\text{dep } f \geq 2$ . We consider the diagonal action of  $G$  on  $\mathfrak{g}^n$  and we have  $J \subset S((\mathfrak{g}^n)^*)^G$ . In section 3, we show how Proposition 1 implies the following Proposition.

---

\*[michael.bulois@univ-angers.fr](mailto:michael.bulois@univ-angers.fr), +33 6 33 12 04 91, LAREMA, 2 boulevard Lavoisier, 49045 Angers CEDEX 1, France.  
Université d'Angers, Laboratoire de Mathématiques d'Angers (LAREMA), UMR 6093.

**Proposition 2.**

$$\mathcal{V}(J) = G.(\mathfrak{b} \times \cdots \times \mathfrak{b}), \quad \mathcal{V}(J_0) = G.(\mathfrak{n} \times \cdots \times \mathfrak{n})$$

where  $\mathfrak{b}$  is any Borel subalgebra of  $\mathfrak{g}$  with nilpotent radical  $\mathfrak{n}$ .

The question of finding such ideals arise naturally when one study the diagonal action of  $G$  on  $\times_{i=1}^n \mathfrak{g}$ . Indeed, when  $n = 1$ ,  $G.\mathfrak{n}$  is the nilpotent cone  $\mathcal{N}$ . In the  $n = 2$  case, several authors pointed out nice generalizations of  $\mathcal{N}$ . Let us mention the set of *nilpotent pairs* of [Gi], whose *principal* elements lie in a finite number of orbits under the diagonal action of  $G$ . Looking at couples of commuting elements, we get the *nilpotent commuting variety*  $\mathfrak{C}^{nil}(\mathfrak{g}) := \{(x, y) \in \mathcal{N} \times \mathcal{N} \mid [x, y] = 0\}$  studied in [Ba, Pr] and which has the nice property of being equidimensional. Finally, the *nilpotent bicone*, studied in [CM], is the affine subscheme  $\mathfrak{N} \subset \mathfrak{g} \times \mathfrak{g}$  defined by the polarized polynomials  $p_i(x + ty) = 0, \forall t \in \mathbb{k}$ . Its underlying set consists of pairs whose any linear combination is nilpotent. It is a non-reduced complete intersection, which contains  $\mathfrak{C}^{nil}(\mathfrak{g})$  and which has  $G.(\mathfrak{n} \times \mathfrak{n})$  has an irreducible component.

*Acknowledgments.* J-Y. Charbonnel drew the author's attention to the plausibility of Proposition 1 and to its importance in view of a characterization of  $G.(\mathfrak{b} \times \mathfrak{b})$  as given in Proposition 2. The author thanks him for these remarks and for his encouragements in the writing of this note.

## 2 Very nilpotent basis

The aim of this section is to prove proposition 1. As a first step, we assume that  $\mathfrak{g}$  is semisimple. We are going to prove that  $\mathfrak{g}$  has no very nilpotent basis, cf. Corollary 5.

First, we have to state some properties of the characteristic grading of a nilpotent element. Let  $y$  be a nilpotent element of  $\mathfrak{g}$  and embed  $y$  in a  $\mathfrak{sl}_2$ -triple  $(y, h, f)$ . Consider the characteristic grading

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}(h, i)$$

where  $\mathfrak{g}(h, i) = \{z \in \mathfrak{g} \mid [h, z] = iz\}$  and denote by  $\text{pr}_i$  the projection  $\mathfrak{g} \rightarrow \mathfrak{g}(h, i)$  with respect to this grading. Then  $\mathfrak{g}(h, 0) = \mathfrak{g}^h$  is a subalgebra of  $\mathfrak{g}$ , reductive in  $\mathfrak{g}$ , i.e.  $\text{ad}_{\mathfrak{g}(h, 0)}(\mathfrak{g})$  is a semisimple representation.

**Lemma 3.** i) An element  $x \in \bigoplus_{i \geq 0} \mathfrak{g}(h, i)$  is nilpotent if and only if  $\text{pr}_0(x)$  is nilpotent in  $\mathfrak{g}(h, 0)$ .  
ii) For all  $i$ ,  $\langle [\mathfrak{g}(h, i), \mathfrak{g}(h, -i)] \rangle \subset \mathfrak{g}(0, h)$  is Lie subalgebra, reductive in  $\mathfrak{g}(0, h)$ .

*Proof.* i) This follows from the fact that  $\bigoplus_{i \geq 0} \mathfrak{g}(h, i)$  is a parabolic subalgebra of  $\mathfrak{g}$  having  $\mathfrak{g}(h, 0)$  as Levi factor.

ii) If  $i = 0$ , the result is straightforward. In the following, we assume  $i \neq 0$ . Embed  $h$  in a Cartan subalgebra  $\mathfrak{h}$ . This gives rise to a root system  $R(\mathfrak{g}, \mathfrak{h}) \subset \mathfrak{h}^*$ . Choose a fundamental basis  $B$  of the root system  $R(\mathfrak{g}, \mathfrak{h})$  such that  $h$  lies in the positive Weyl Chamber associated to  $B$ .

Let  $[x_1, y_1], [x_2, y_2]$  be two elements of  $[\mathfrak{g}(h, i), \mathfrak{g}(h, -i)]$ . Then,

$$[[x_1, y_1], [x_2, y_2]] = \left[ [[x_1, y_1], x_2], y_2 \right] + \left[ x_2, [[x_1, y_1], y_2] \right].$$

Since  $x_2$  and  $[[x_1, y_1], x_2]$  (resp.  $y_2$  and  $[[x_1, y_1], y_2]$ ) are elements of  $\mathfrak{g}(h, i)$  (resp.  $\mathfrak{g}(h, -i)$ ), the element  $[[x_1, y_1], [x_2, y_2]]$  belongs to  $\langle [\mathfrak{g}(h, i), \mathfrak{g}(h, -i)] \rangle$ . By linearity, we deduce that  $\langle [\mathfrak{g}(h, i), \mathfrak{g}(h, -i)] \rangle$  is a Lie subalgebra of  $\mathfrak{g}(0, h)$ .

Write  $R_i := \{\alpha \in R(\mathfrak{g}, \mathfrak{h}) \mid \alpha(h) = i\}$ . Hence  $\mathfrak{g}(h, i) = \bigoplus_{\alpha \in R_i} \mathfrak{g}^\alpha$ , where  $\mathfrak{g}^\alpha$  is the root space associated to  $\alpha$ . Then, we see that

$$\bigoplus_{R_{i, -i}} \mathfrak{g}^\alpha \subset \langle [\mathfrak{g}(h, i), \mathfrak{g}(h, -i)] \rangle \subset \mathfrak{h} \oplus \bigoplus_{R_{i, -i}} \mathfrak{g}^\alpha, \quad (1)$$

where  $R_{i,-i} = (R_i + R_{-i}) \cap R(\mathfrak{g}, \mathfrak{h}) = (R_i - R_i) \cap R(\mathfrak{g}, \mathfrak{h})$ . Since  $R_{i,-i} = -R_{i,-i}$ ,  $\langle [\mathfrak{g}(h, i), \mathfrak{g}(h, -i)] \rangle$  is reductive and it follows from (1) that its central elements are semisimple in  $\mathfrak{g}$ . Hence the result.  $\square$

The main step of the proof of Proposition 1 lies in the following lemma.

**Lemma 4.** *If  $\mathfrak{g}$  has a very nilpotent basis, then there exists a non-zero proper Lie subalgebra  $\mathfrak{k} \subset \mathfrak{g}$ , reductive in  $\mathfrak{g}$ , and a basis  $(x_1, \dots, x_p)$  of  $\mathfrak{k}$  such that  $\text{ad}_z$  is nilpotent in  $\mathfrak{gl}(\mathfrak{g})$  for each  $z \in \mathcal{I}(x_1, \dots, x_p)$ .*

*Proof.* Let  $(y_1, \dots, y_n)$  be a very nilpotent basis of  $\mathfrak{g}$ . Each  $y_i$  is nilpotent. Embed  $y_1$  in a  $\mathfrak{sl}_2$ -triple  $(y_1, h, f)$  and consider the characteristic grading

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}(h, i).$$

Denote by  $i_0$  the highest weight in this decomposition, i.e.  $\mathfrak{g}(h, i_0) \neq \{0\}$  and  $\mathfrak{g}(h, i) = \{0\}$  for all  $i > i_0$ . We set  $\mathfrak{k} := \langle [\mathfrak{g}(h, i_0), \mathfrak{g}(h, -i_0)] \rangle$ . Since  $\mathfrak{k} \subset \mathfrak{g}(0, h) \neq \mathfrak{g}$  is reductive in  $\mathfrak{g}$  (Lemma 3), there remains to find the basis  $(x_1, \dots, x_p)$ .

The endomorphism  $\text{ad}_{y_1}$  is nilpotent of order  $i_0 + 1$  and

$$(\text{ad}_{y_1})^{i_0} : \mathfrak{g} \rightarrow \mathfrak{g}(h, i_0)$$

is surjective. We define  $z_j := \text{ad}_{y_1}^{i_0}(y_j)$  for  $j \in J := \llbracket 1, n \rrbracket$ . By construction,  $z_j \in \mathcal{I}(y_1, \dots, y_n)$  and  $(z_j)_{j \in J}$  is a family spanning the vector space  $\mathfrak{g}(h, i_0)$ . On the opposite side, we define  $y'_k := \text{pr}_{-i_0}(y_k)$  for  $k \in J$ . The family  $(y'_k)_{k \in J}$  spans the vector space  $\mathfrak{g}(h, -i_0)$ .

Consider now  $\text{ad}_{z_j} : \mathfrak{g} \rightarrow \bigoplus_{i \geq 0} \mathfrak{g}(h, i)$  and define

$$x_{j,k} := \text{pr}_0 \circ \text{ad}_{z_j}(y_k) = \text{ad}_{z_j} \circ \text{pr}_{-i_0}(y_k), \quad j, k \in J^2.$$

The family  $(x_{j,k})_{j,k}$  spans the vector space  $\mathfrak{k} \subset \mathfrak{g}(h, 0)$ . The elements  $\text{ad}_{z_j}(y_k)$  belong to  $\mathcal{I}(y_1, \dots, y_n)$ . Hence, it follows from lemma 3 that the elements  $x_{j,k}$  and there iterated brackets are nilpotent. In other words, if  $(x_1, \dots, x_p)$  is a basis of  $\mathfrak{k}$  extracted from  $(x_{j,k})_{j,k \in J^2}$ , then it fulfills the required properties.  $\square$

**Corollary 5.** *Let  $\mathfrak{g} \neq \{0\}$  be a semisimple Lie algebra, then  $\mathfrak{g}$  has no very nilpotent basis*

*Proof.* We argue by induction on  $\dim \mathfrak{g}$ . Assume that there is no semisimple  $\mathfrak{w} \subset \mathfrak{g}$  such that  $0 \neq \dim \mathfrak{w} < \dim \mathfrak{g}$  having a very nilpotent basis. Assume that  $\mathfrak{g}$  has one. Then, we define the reductive subalgebra  $\mathfrak{k}$  equipped with the basis  $(x_1, \dots, x_p)$  as in Lemma 4. By hypothesis  $\mathfrak{k}$  is not semisimple. Hence  $\mathfrak{k}$  has a non-trivial centre whose elements are semisimple. Therefore there must be some  $i \in \llbracket 1, p \rrbracket$  such that  $x_i$  is not nilpotent and we get a contradiction.  $\square$

We are now ready to finish the **proof of proposition 1** in the general case. From now on, we forget the semisimplicity assumption on  $\mathfrak{g}$ .

First of all, we note that whenever  $\mathfrak{g}$  is nilpotent, then any basis of  $\mathfrak{g}$  is very nilpotent.

Conversely, assume that  $\mathfrak{g}$  is any Lie algebra having a very nilpotent basis  $(y_1, \dots, y_n)$ . Let  $\mathfrak{r}$  be the radical of  $\mathfrak{g}$ . The algebra  $\mathfrak{g}/\mathfrak{r}$  is semisimple and we can extract a very nilpotent basis  $(x_1, \dots, x_p)$  of  $\mathfrak{g}/\mathfrak{r}$  from the projection of the elements  $y_i$  via  $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{r}$ . It follows from corollary 5 that  $\mathfrak{g}/\mathfrak{r} = \{0\}$ . In other words  $\mathfrak{g}$  is solvable.

Then, one may apply Lie's Theorem (see, e.g., [TY, 19.4.4]). It states that  $\text{ad}_{\mathfrak{g}} \subset \mathfrak{gl}(\mathfrak{g})$  can be seen as a subspace of a set of upper triangular matrices. In fact, the nilpotency condition on the  $\text{ad}_{y_i}$  implies that  $\text{ad}_{\mathfrak{g}}$  is a subspace of a set of strictly upper triangular matrices. In particular,  $\mathfrak{g}$  is nilpotent. This ends the proof of Proposition 1.

### 3 $n$ -tuples lying in a same Borel subalgebra

In this section  $\mathfrak{g}$  is assumed to be semisimple. Let  $\mathfrak{b}$  be a Borel subalgebra of  $\mathfrak{g}$  and  $\mathfrak{n}$  be the nilradical of  $\mathfrak{b}$ . Define two ideals of  $S((\mathfrak{g}^n)^*)^G$  by:

$$J_0 := (p_i \circ f \mid i \in \llbracket 1, d \rrbracket, f \in \mathcal{I}_n), \quad J := (p_i \circ f \mid i \in \llbracket 1, d \rrbracket, f \in \mathcal{I}_n, \text{dep } f \geq 2),$$

where  $p_1, \dots, p_d$  are as in the introduction.

**Proposition 6.**

$$\begin{aligned} \mathcal{V}(J_0) &= G.(\mathfrak{n} \times \dots \times \mathfrak{n}), \\ \mathcal{V}(J) &= G.(\mathfrak{b} \times \dots \times \mathfrak{b}). \end{aligned}$$

*Proof.* The inclusions  $\mathcal{V}(J_0) \supset G.(\mathfrak{n} \times \dots \times \mathfrak{n})$  and  $\mathcal{V}(J) \supset G.(\mathfrak{b} \times \dots \times \mathfrak{b})$  are straightforward.

Let us prove the reverse inclusions. Let  $(y_1, \dots, y_n) \in \mathfrak{g}$  and let  $\mathfrak{k} \subset \mathfrak{g}$  be the Lie subalgebra generated by the elements  $(y_1, \dots, y_n)$ . Assume that  $(y_1, \dots, y_n) \in \mathcal{V}(J_0)$ . Choose a basis of  $\mathfrak{k}$ ,  $(z_1, \dots, z_p) \in (\mathcal{I}(y_1, \dots, y_n))^p$ . Then  $(z_1, \dots, z_p)$  is a very nilpotent basis of  $\mathfrak{k}$ . It follows from Proposition 1 that  $\mathfrak{k}$  is nilpotent. Hence there exists  $g \in G$  such that  $g.\mathfrak{n} \supset \mathfrak{k}$  and  $(y_1, \dots, y_n) \in g.(\mathfrak{n} \times \dots \times \mathfrak{n})$ .

Assume now that  $(y_1, \dots, y_n) \in \mathcal{V}(J)$ . Arguing along the same lines, one finds that  $[\mathfrak{k}, \mathfrak{k}]$  is nilpotent. Hence  $\mathfrak{k}$  is solvable and there exists  $g \in G$  such that  $g.\mathfrak{b} \supset \mathfrak{k}$ .  $\square$

The ideals  $J_0$  and  $J$  are defined by making use of an infinite number of generators. In fact, if one is more careful with the arguments of section 2, it is possible to restrict to a finite number. Let us sketch the proof of this. A rough estimation shows that, for  $(y_1, \dots, y_n) \in \mathfrak{g}^n$ , the subalgebra  $\mathfrak{k}$  generated by  $(y_1, \dots, y_n)$  is spanned by  $\{f(y_1, \dots, y_n) \mid f \in \mathcal{I}_n, \text{dep } f \leq \dim \mathfrak{g}\}$  as a vector space. Then, assume that  $(z_1, \dots, z_p)$  is a basis of a semisimple Lie algebra  $\mathfrak{k}$ . One can restrict in the proof of Lemma 4 and Corollary 5 to the assumption that  $f(z_1, \dots, z_p)$  is nilpotent for  $\text{dep } f \leq \dim \mathfrak{k}$ . Defining

$$J_0 := (p_i \circ f \mid \text{dep } f \leq (\dim \mathfrak{g})^2), \quad J := (p_i \circ f \mid 2 \leq \text{dep } f \leq 2(\dim \mathfrak{g})^2),$$

we claim that:

**Claim 7.**  $\mathcal{V}(J') = \mathcal{V}(J)$  and  $\mathcal{V}(J'_0) = \mathcal{V}(J_0)$ .

## References

- [Ba] V. Baranovsky, The variety of pairs of commuting nilpotent matrices is irreducible, *Transform. Groups* **6** (2001), 3-8.
- [CM] J. Y. Charbonnel and A. Moreau, Nilpotent bicone and characteristic submodule in a reductive Lie algebra, *Transform. Groups* **14** (2009), 319-360.
- [Gi] V. Ginzburg, Principal nilpotent pairs in a semisimple Lie algebra I., *Invent. Math.* **140** (2000), 511-561.
- [Pr] A. Premet, Nilpotent commuting varieties of reductive Lie algebras, *Invent. Math.* **154** (2003), 653-683.
- [TY] P. Tauvel and R. W. T. Yu, *Lie algebras and algebraic groups*, Springer Monographs in Mathematics, Springer-Verlag, 2005.